

Physics 121. Lecture 09.



F/A-18 Hornet breaking the sound barrier. Photo by John Gay

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

Navy Lt. Ron Candiloro's F/A-18 Hornet creates a shock wave as he breaks the sound barrier July 7. The shock wave is visible as a large cloud of condensation formed by the cooling of the air. A smaller shock wave can be seen forming on top of the canopy. It is possible for a skilled pilot to work the plane's throttle to move the shock wave forward or aft.

1

Physics 121. Lecture 09.

• Topics:

- Course announcements
- Quiz
- Work and Energy:
 - Definition
 - Work done by a Variable Force
 - Kinetic Energy

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

2

Physics 121. Course announcements.

- Homework set # 4 is due on February 28 at 8.30 am.
 - Note there is written component on this assignment.
 - The most effective way to work on the assignments is to tackle 1 or 2 problems a day.
- On Monday 2/23, you will receive an email with your results of Exam # 1.
- Exam # 1 will be returned during workshops next week.
 - The minimum you should do is to compare the scores shown on your blue booklet with the scores in my email.
 - You should of course compare your solutions with the posted solutions.
 - If you are not happy with the grading of the exam, you need to write down why you feel you deserve more points on a particular question, attach the note to your blue booklet(s), and return them to me before the end of lecture 13 (March 5).

Frank L. H. Wolfs


Department of Physics and Astronomy, University of Rochester

3

Some observations on Exam # 1.

13% did not (correctly) enter the ID #.

- **Scantron:**
 - 14 students did not enter their student ID on the scantron form.
 - 9 students did not enter their student ID correctly.
 - Students entered their solutions incorrectly.
 - Answers not on front.
- This time, I added missing student IDs were possible, but **there will be no adjustments for Exam # 2.** But there were forms where I could not even read the name to determine the student ID.
- Note: your ID has 8 numbers.




Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester

4

First results Exam # 1.

Q1 – Q10.



Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester

5

Quiz.

- Let's practice what we learned (and read).



Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester

6

Work and energy.

- In the next few weeks, we will not discuss any new physics, but develop tools to simplify how we use our understanding of the force laws and the laws of motion to understand and/or predict the outcome of experiments.
- We will start with defining the concept of work and the concept of energy.

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester


7

Work and energy.

- When a force F is applied to an object, it may produce a displacement d .
- The work W done by the force F is defined as

$$W = \vec{F} \cdot \vec{d} = Fd\cos\theta$$

where θ is the angle between the force $\vec{F}$ and the displacement $\vec{d}$.



Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester


8

Work.

- Consider the definition of W :

$$W = \vec{F} \cdot \vec{d} = Fd\cos\theta$$

- The work done by the force $\vec{F}$ is zero if:
 - $d = 0$ m (no displacement)
 - $\theta = 90^\circ$ (force perpendicular to the displacement).



Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

9

Work.

Positive, zero, or negative.

- Work done by a force can be positive, zero, or negative, depending on the angle θ :
- When $0^\circ \leq \theta < 90^\circ$ (scalar product between F and d is larger than 0) the speed of the object will increase.
- When $\theta = 90^\circ$ (scalar product between F and d is 0) the speed of the object will not change.
- When $90^\circ < \theta \leq 180^\circ$ (scalar product between F and d is smaller than 0) the speed of the object will decrease.

Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester

10

Work: units.

- The unit of work is the Joule (abbreviated J).
- Per definition, $1 \text{ J} = 1 \text{ Nm} = 1 \text{ kg m}^2/\text{s}^2$.
- There are many important examples of forces that do not do any work. For example, the gravitational force between the earth and the moon does not do any work! Note: in this case, the speed of the moon does not change.

Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester

11

Power.

- In many cases, the work done by a tool is less important than the rate with which the work can be done.
- For example, explosive devices get their properties from being able to do a lot of work over a very short period in time. The same amount of work done over a longer period of time might not lead to destruction.

Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester

12

Power: units.

- Power is defined as work per unit time:

$$P = \frac{dW}{dt}$$


- The unit of power is the Watt, abbreviated by a W.
- Per definition:

$$1 \text{ W} = 1 \text{ J/s} = 1 \text{ kg m}^2/\text{s}^3$$

- The power you consume at home is often expressed in terms of kWh, which is the use of 1 kW of power

for 1 hour.
Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester




13

Work. An example problem.

A block of mass M is drawn at constant speed a distance d along a horizontal floor by a rope exerting a force $\vec{F}$ at an angle θ above the horizontal. Compute

- the work done by the rope on the block.
- the coefficient of kinetic friction between block and floor.



Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

14

Work. An example problem: step 1.

- The requirement that the object moved with constant speed tells us that the net force acting on it must be 0 N.
- Thus, the net forces in the x and y directions:

$$\Sigma F_x = F \cos \theta - f_k$$

$$\Sigma F_y = N + F \sin \theta - Mg$$

..... must be zero.

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

15


Work.

An example problem: step 2.

- The normal force N can be determined based on the fact that the net force in the vertical direction must be zero:

$$N = Mg - F \sin \theta$$

- Based on the now known normal force N we can determine the frictional force f (kinetic friction since the block is moving):

$$f_k = \mu_k N = \mu_k (Mg - F \sin \theta)$$


Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester

16


Work.

An example problem: step 3.

- The friction force is also directly related to the applied force F by considering net force in the horizontal direction, which has to be zero.
- This requires that

$$f_k = F \cos \theta$$

- The work done by the friction force is negative (since direction and displacement are in opposite directions):

$$W_f = \vec{f}_k \cdot \vec{d} = -f_k d = -F d \cos \theta$$


Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester

17

Work.

An example problem: step 4.


- The work done by the applied force F is equal to

$$W_F = \vec{F} \cdot \vec{d} = F d \cos \theta$$

- The net work done (due to the applied force and the friction force) is equal to zero.

$$W_{total} = \sum W_i = W_F + W_f = 0$$

- This is not really a surprise since the net force on the object is equal to zero.



Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester

18

4 Minute 9 Second Intermission



- Since paying attention for 1 hour and 15 minutes is hard when the topic is physics, let's take a 4 minute 9 second intermission.
- You can:
 - Stretch out.
 - Talk to your neighbors.
 - Ask me a quick question.
 - Enjoy the fantastic music.
 - Solve a WeBWork problem.
 - Go asleep, as long as you wake up in 4 minutes and 9 seconds.




Frank L. H. Wolfs
Department of Physics and Astronomy, University of Rochester

19

Work done by a varying force.


- In most realistic cases, we need to consider the work done when the force is varying (both in magnitude and direction) as function of time and/or position.
- In this case, we can still use the same approach as we just discussed by breaking up the motion into small intervals such that the path is linear and the force is constant during the intervals considered.



Frank L. H. Wolfs
Department of Physics and Astronomy, University of Rochester

20

Work done by a varying force.



Frank L. H. Wolfs
Department of Physics and Astronomy, University of Rochester

21


Calculating work. Do less work by thinking before starting!

- Consider the work done by all forces acting on the pendulum when it moves from position 1 to position 2.
- During this motion, the angle between the path and the net force changes. What am I to do?

$$W = \vec{F}_t \cdot \vec{d} = (\vec{F} + \vec{F}_g) \cdot \vec{d}$$

$$\vec{F} \cdot \vec{d} = F_x = F \sqrt{h(2r-h)}$$

$$\vec{F}_g \cdot \vec{d} = -F_g h = -mgh$$



Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

22

The work-energy theorem.




- We have already seen that there is a connection between the work done by a force and the change in the speed of the object:
 - If $W > 0$ J: speed increases
 - If $W = 0$ J: speed remains constant
 - If $W < 0$ J: speed decreases

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

23

The work-energy theorem.



- Consider the bus starting from rest ($v_1 = 0$ m/s) and having an acceleration $a = F/m$ (assume $v \ll c$). The velocity at a later time t will be equal to

$$v(t) = v_0 + at = at$$
- This relation can be used to determine the time t at which the bus reaches a certain velocity v_2 :


$$t = \frac{v_2}{a}$$

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

24

The work-energy theorem.



- The displacement at this time t is equal to

$$d = x\left(t = \frac{v_2}{a}\right) = x_0 + v_0 t + \frac{1}{2} a t^2 = \frac{1}{2} a \left(\frac{v_2}{a}\right)^2 = \frac{1}{2} \frac{v_2^2}{a}$$

- The work done by the force F during this period is equal to


$$W = \vec{F} \cdot \vec{d} = m a \left(\frac{1}{2} \frac{v_2^2}{a}\right) = \frac{1}{2} m v_2^2 \leftarrow \text{Kinetic energy } K$$

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

25

The work-energy theorem.



- Assuming that there is no heat flow in or out of the bus, we conclude that for low velocities (non-relativistic velocities):

The net work done on an object is equal to the change in its kinetic energy.

- In the case of the bus: $F_{net} d = \Delta E = \frac{1}{2} m v_2^2 - \frac{1}{2} m v_1^2$

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester


26

The energy principle. Example 1.

- An object with mass m is at rest at time $t = 0$ s. It falls under the influence of gravity through a distance h . What is its velocity at that point?

- Solution:

- Assume $Q = 0$.
- Work done by the gravitational force = mgh .
- Change in energy = change in kinetic energy = $(1/2)mv_1^2$.
- The Energy Principle ($Q = 0$): $mgh = (1/2)mv_1^2$ or $v_1 = \sqrt{2gh}$



Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

27

The work-energy theorem.

An application.

- An object with mass m is at rest at time $t = 0$ s. It falls under the influence of gravity through a distance h . What is its velocity at that point?
- Solution:
 - Initial kinetic energy: $K_i = 0$ J.
 - Work done by the gravitational force: $W = mgh$.
 - Change in kinetic energy: $K_f - K_i = \frac{1}{2}mv_f^2$.
 - Work-Energy theorem:

$mgh = \frac{1}{2}mv_f^2$

or

$v_f = \sqrt{2gh}$

y-axis

y0 = 0 m

v0 = 0 m/s

W

y1 = -h

v1 = ?

W


Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

28

That's all!

Next week: conservation of energy.



Unusual Spherules on Mars

Credit: [Mars Exploration Rover Mission](#), [JPL](#), [USGS](#), [NASA](#)

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

29
