

Physics 121.
Lecture 17.



Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 1

1

Physics 121.
Lecture 17.

- Course Information
- Topics to be discussed today:
 - Review of Rotational Motion
 - Rolling Motion
 - Angular Momentum
 - Conservation of Angular Momentum

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 2

2

Physics 121.
Lecture 17.


- Homework set # 6 is now available and is due on Saturday April 4 at 8.30 am.
- The grades for exam # 2 were distributed via email on Saturday March 28.
- You should pick up your exam in workshop this week. Please check it carefully for any errors.
- If you are not happy with the grading of your exam, you need to write down why you feel you deserve more points on a particular question, attach the note to your blue booklet(s), and return them to me before the end of lecture 20 (April 9).

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 3

3

Results of individual questions on Exam # 2.




Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 4

4

Grade Distribution Exam # 2.




Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 5

5


Your prediction versus reality.
Reality is better than your prediction.




Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 6


6



7



8



9


Rotational variables. A quick review.

- The variables that are used to describe rotational motion are:

- Angular position θ
- Angular velocity $\omega = d\theta/dt$
- Angular acceleration $\alpha = d\omega/dt$

- The rotational variables are related to the linear variables:

- Linear position $l = R\theta$
- Linear velocity $v = R\omega$
- Linear acceleration $a = R\alpha$



Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester, Slide 10

10

Rotational kinetic energy. A quick review.

- Since the components of a rotating object have a non-zero (linear) velocity we can associate a kinetic energy with the rotational motion:

$$K = \sum_i \frac{1}{2} m_i v_i^2 = \sum_i \frac{1}{2} m_i (\omega r_i)^2 = \frac{1}{2} (\sum_i m_i r_i^2) \omega^2$$

- The kinetic energy is proportional to the square of the rotational velocity ω . Note: the equation is similar to the translational kinetic energy ($\frac{1}{2} m v^2$) except that instead of being proportional to the mass m of the object, the rotational kinetic energy is proportional to the **moment of inertia I** of the object:

$$I = \sum_i m_i r_i^2 \Rightarrow K = \frac{1}{2} I \omega^2$$

Note: units of I : kg m^2

Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester, Slide 11


11

Torque. A quick review.

- In general, the torque associated with a force F is equal to

$$|\vec{\tau}| = r F \sin\phi = |\vec{r} \times \vec{F}|$$

- The arm of the force (also called the moment arm) is defined as $r \sin\phi$. The arm of the force is the perpendicular distance of the axis of rotation from the line of action of the force.
- If the arm of the force is 0, the torque is 0, and there will be no rotation.
- The maximum torque is achieved when the angle ϕ is 90° .



Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester, Slide 12

12

Torque. A quick review.

- The torque τ of the force F is related to the angular acceleration α :


$$\vec{\tau} = I\vec{\alpha}$$

- This equation looks similar to Newton's second law for linear motion:

$$\vec{F} = m\vec{a}$$

- Corresponding variables for linear and rotational motion:

linear	rotational
mass m	moment I
force $\vec{F}$	torque $\vec{\tau}$



The direction of the torque defines the direction of the angular acceleration.

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 13

13

Rolling motion. A quick review.

- Rolling motion is a combination of translational and rotational motion.

- The kinetic energy of rolling motion has thus two contributions:

- Translational kinetic energy:


$$\frac{1}{2}Mv_{CM}^2$$

- Rotational kinetic energy:

$$\frac{1}{2}I_{CM}\omega^2$$

- Assuming the wheel does not slip:

$$\omega = \frac{v}{R}$$



Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 14

14

How different is a world with rotational motion? Sample problem.

- Consider the loop-to-loop. What height h is required to make it to the top of the loop?

- First consider the case without rotation:

- Initial mechanical energy: mgh .


- Minimum velocity at the top of the loop is determined by requiring that $mv^2/R > mg$ or $v^2 > gR$.

- The mechanical energy at the top of the loop is thus equal to

$$\frac{1}{2}mv^2 + 2mgR > \frac{5}{2}mgR$$

- Conservation of energy requires

$$h > \frac{5}{2}R$$




Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 15

15

How different is a world with rotational motion? Sample problem.

- What changes when the object rotates?
 - The minimum velocity at the top of the loop will not change.
 - The minimum translational kinetic energy at the top of the loop will not change.
 - But in addition to translational kinetic energy, there is now also rotational kinetic energy.
 - The minimum mechanical energy is at the top of the loop has thus increased.
 - The required minimum height must thus have increased.
- OK, let's now calculate by how much the minimum height has increased.



Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester, Slide 16

16

How different is a world with rotational motion? Sample problem.

- The total kinetic energy at the top of the loop is equal to

$$K_f = \frac{1}{2}I\omega^2 + \frac{1}{2}Mv^2 = \frac{1}{2}\left(\frac{I}{R^2} + M\right)v^2$$


- This expression can be rewritten as

$$K_f = \frac{1}{2}\left(\frac{2}{5}M + M\right)v^2 = \frac{7}{10}Mv^2$$

- This is the minimum mechanical energy required to reach this point, and we can determine the minimum height:

$$h \geq \frac{27}{10}R$$

Note: without rotation $h \geq 25/10 R$!!!



Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester, Slide 17

17



3 Minute 52 Second Intermission.

- Since paying attention for 1 hour and 15 minutes is hard when the topic is physics, let's take a 3 minute 52 second intermission.

- You can:
 - Stretch out.
 - Talk to your neighbors.
 - Ask me a quick question.
 - Enjoy the fantastic music.



Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester, Slide 18

18

Quiz.

- Let us practice what we have learned.




Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 19

19

Torque.

- The torque associated with a force is a vector. It has a magnitude and a direction.
- The direction of the torque can be found by using the right-hand rule to evaluate $\vec{r} \times \vec{F}$.
- For extended objects, the total torque is equal to the vector sum of the torque associated with each "component" of this object.



Frank L. H. Wolfs


Department of Physics and Astronomy, University of Rochester, Slide 20

20

Angular momentum.

- We have seen many similarities between the way in which we describe linear and rotational motion.
- Our treatment of these types of motion are similar if we recognize the following equivalence:

linear	rotational
mass m	moment I
force $\vec{F}$	torque $\vec{\tau} = \vec{r} \times \vec{F}$



- What is the equivalent to linear momentum?
Answer: angular momentum.

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 21


21

Angular momentum.

- The **angular momentum** is defined as the vector product between the position vector and the linear momentum.

Note:

- Compare this definition with the definition of the torque.
- Angular momentum is a vector.
- The unit of angular momentum is $\text{kg m}^2/\text{s}$.
- The angular momentum depends on both the magnitude and the direction of the position and linear momentum vectors.
- Under certain circumstances the angular momentum of a system is conserved!



Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 22


22

Angular momentum.

- Consider an object carrying out circular motion.
- For this type of motion, the position vector will be perpendicular to the momentum vector.
- The magnitude of the angular momentum is equal to the product of the magnitude of the radius $\vec{r}$ and the linear momentum $\vec{p}$:

$$L = mvr = mr^2 \frac{v}{r} = I\omega$$

- Note: compare this with the linear momentum $p = mv$!



Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 23


23

Angular momentum.

- An object does not need to carry out rotational motion to have an angular momentum.
- Consider a particle P carrying out linear motion in the xy plane.
- The angular momentum of P (with respect to the origin) is equal to

$$\vec{L} = \vec{r} \times \vec{p} = mrv \sin \theta \hat{k} = pr_{\perp} \hat{k}$$

and will be constant (if the linear momentum is constant).



Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 24

24

Conservation of angular momentum.

- Consider the change in the angular momentum of a particle:

$$\frac{d\vec{L}}{dt} = \frac{d}{dt}(\vec{r} \times \vec{p}) = m \left(\vec{r} \times \frac{d\vec{v}}{dt} + \frac{d\vec{r}}{dt} \times \vec{v} \right) = m(\vec{r} \times \vec{a} + \vec{v} \times \vec{v})$$

$$\frac{d\vec{L}}{dt} = \vec{r} \times m\vec{a} = \vec{r} \times \Sigma \vec{F} = \Sigma \vec{\tau}$$

- Consider what happens when the net torque is equal to 0:

$$\frac{d\vec{L}}{dt} = 0 \Rightarrow L = \text{constant}$$

- When we take the sum of all torques, the torques due to the internal forces cancel and the sum is equal to torque due to all external forces.
- Note: notice again the similarities between linear and rotational motion.

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 25

25


Conservation of angular momentum.

- The connection between the angular momentum $\vec{L}$ and the torque $\vec{\tau}$

$$\Sigma \vec{\tau} = \frac{d\vec{L}}{dt}$$

is only true if $\vec{L}$ and $\vec{\tau}$ are calculated with respect to the same reference point (which is at rest in an inertial reference frame).

- The relation is also true if $\vec{L}$ and $\vec{\tau}$ are calculated with respect to the center of mass of the object (note: center of mass can accelerate).



Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 26

26

Conservation of angular momentum.

- Ignoring the mass of the bicycle wheel, the external torque will be close to zero if we use the center of the disk as our reference point.

- Since the external torque is zero, angular momentum thus should be conserved.

- I can change the orientation of the wheel by applying internal forces. In which direction will I need to spin to conserve angular momentum?



Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 27

27

Quiz and concept test.

- Let us practice what we have learned.



Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 28

28

Done for today!
More about rotational motion on Thursday.



The Orion Nebula from CFHT

Credit & [Copyright: Canada-France-Hawaii Telescope, J.-C. Cuillandre \(CFHT\), Caelum](#)

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 29

29
