

Classical Mechanics
Phy 235, Lecture 24.

Frank L. H. Wolfs
Department of Physics and Astronomy
University of Rochester

Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester, Slide 1

1


PH-AKA arriving in Montreal.



Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester, Slide 2

2

Friday December 5.
An important evening: pakjes avond.



Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester, Slide 3

3


Announcements

- Homework set # 10 is due on Friday December 5 at noon.
- On December 1, the University opened the end-of-semester course survey.
 - The survey closes on December 11.
 - I encourage all of you to complete this survey. Your opinion matters!
 - If by December 11, 95% of the PHY 235 students have completed the survey, I add 5% to everyone's final exam score.
- There will be no recitations next week, but office hours will be continue as scheduled to help you with final-exam related questions.
- The final exam will take place on Sunday December 14 between 12.30 pm and 3.30 pm in Dewey 1101.

Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester, Slide 4

4

Chapter 13: from the loaded string to a “real” string.



Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester, Slide 5

5

Chapter 13: from the loaded string to a “real” string.

- Consider the following:
 - Increase the number of masses n to infinity.
 - Decrease the distance d to zero such that $(n + 1)d = L$.
 - Decrease the mass m to zero such that $m/d = \text{constant}$.
- In this case, the displacement q can be written as:

$$q_j(t) = \sum_s \beta_s \sin\left(s\pi \frac{x}{L}\right) e^{i\omega_s t} = q(x, t)$$
- The eigen frequencies are

$$\omega_s = \frac{s\pi}{L} \sqrt{\frac{\tau}{\rho}}$$

Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester, Slide 6

6

Energy in a string.

- Kinetic energy:

$$T = \int_0^L dT = \frac{L}{4} \rho \sum_s (\omega_s v_s \cos(\omega_s t) + \omega_s \mu_s \sin(\omega_s t))^2$$

- Potential energy:

$$U = \frac{\tau}{2} \int_0^L \left(\frac{\partial q}{\partial x} \right)^2 dx = \frac{\rho L}{4} \sum_s \omega_s^2 (\mu_s \cos(\omega_s t) - v_s \sin(\omega_s t))^2$$

- Total energy:


$$E = T + U = \frac{\rho L}{4} \sum_s \omega_s^2 (\mu_s^2 + v_s^2) = \text{constant}$$

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 7

7

The wave equation.



Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 8

8

The wave equation.

- The ideal wave equation:

$$\frac{\partial^2 q}{\partial x^2} = \frac{\rho}{\tau} \frac{\partial^2 q}{\partial t^2}$$

- The "real" wave equation:

$$\frac{\partial^2 q}{\partial x^2} - \frac{D}{\tau} \frac{\partial q}{\partial t} + \frac{F(x,t)}{\tau} = \frac{\rho}{\tau} \frac{\partial^2 q}{\partial t^2}$$

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 9

9



2 Minute 27 Second Intermission.

- Since paying attention for 1 hour and 15 minutes is hard when the topic is physics, let's take a 2 minute 27 second intermission.
- You can:
 - Stretch out.
 - Talk to your neighbors.
 - Ask me a quick question.
 - Enjoy the fantastic music.



Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 10

10

Problem 13.11.

- When a particular driving force is applied to a string, it is observed that the string's vibration is purely in the n^{th} harmonic. Find the driving force.

$$f_s(t) = \int_0^L F(x,t) \sin \frac{s\pi x}{b} dx = 0 \quad \text{for } s \neq n$$

$$= \int_0^L F(x,t) \sin \frac{s\pi x}{b} dx \neq 0 \quad \text{for } s = n$$

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 11

11

Solving the ideal wave equation.

- The ideal wave equation:

$$\frac{\partial^2 q}{\partial x^2} = \frac{\rho}{\tau} \frac{\partial^2 q}{\partial t^2} = \frac{1}{v^2} \frac{\partial^2 q}{\partial t^2}$$
- No dissipation: energy is conserved.
- Use **separation of variables** to solve the wave equation:

$$q(x,t) = \psi(x)\chi(t)$$

- This results on two differential equations:

$$\frac{v^2 \partial^2 \psi}{\psi \partial x^2} = \frac{1 \partial^2 \chi}{\chi \partial t^2} = \text{constant} = \omega^2$$

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 12

12

Solving the ideal wave equation.

$$\frac{\partial^2 \psi}{\partial x^2} - \frac{\omega^2}{v^2} \psi = 0$$
$$\frac{\partial^2 \chi}{\partial t^2} - \omega^2 \chi = 0$$

Solutions:

$$\psi(x) = e^{\pm i \left(\frac{\omega}{v}\right) x}$$
$$\chi(t) = e^{\pm i \omega t}$$

Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester, Slide 13

13

ENOUGH FOR TODAY?

Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester, Slide 14

14
